

A note on the non-existence of σ -model solitons in the $2 + 1$ dimensional AdS gravity

Piotr Bizoń* and Arthur Wasserman†

**Institute of Physics, Jagellonian University, Kraków, Poland*

†Department of Mathematics, University of Michigan, Ann Arbor, Michigan

February 7, 2008

Abstract

We show that the gravitating static soliton in the $2 + 1$ dimensional $O(3)$ σ model does not exist in the presence of a negative cosmological constant.

1 Introduction

The $2 + 1$ dimensional $O(3)$ σ model coupled to gravity is a wave map $X : M \rightarrow N$ from a $2 + 1$ dimensional spacetime (M, g_{ab}) into a two-sphere S^2 with the round metric G_{AB} defined by the action

$$S = \int_M \left(\frac{R + 2\Lambda}{16\pi G} + L_{WM} \right) dv_g \quad (1)$$

with the Lagrangian density

$$L_{WM} = -\frac{f_\pi^2}{2} g^{ab} \partial_a X^A \partial_b X^B G_{AB}. \quad (2)$$

Here Λ is a cosmological constant, G is the Newton constant and f_π^2 is the wave map coupling constant. The product $\alpha = 8\pi G f_\pi^2$ is dimensionless. The field equations derived from (1) are the wave map equation

$$\square_g X^A + \Gamma_{BC}^A(X) \partial_a X^B \partial_b X^C g^{ab} = 0, \quad (3)$$

where $\Gamma_{BC}^A(X)$ are the Christoffel symbols of the target metric G_{AB} and $\square_g$ is the wave operator associated with the metric g_{ab} , and the Einstein equations $R_{ab} - \frac{1}{2}g_{ab}R + \Lambda g_{ab} = 8\pi G T_{ab}$ with the stress-energy tensor

$$T_{ab} = f_\pi^2 \left(\partial_a X^A \partial_b X^B - \frac{1}{2} g_{ab} (g^{cd} \partial_c X^A \partial_d X^B) \right) G_{AB}. \quad (4)$$

In polar coordinates $X^A = (F, \Phi)$ on the target S^2 the metric takes the form

$$G_{AB}dX^AdX^B = dF^2 + \sin^2 F d\Phi^2. \quad (5)$$

For the domain manifold we take a spherically symmetric 2 + 1 dimensional spacetime and parametrize the metric using areal coordinates

$$g_{ab}dx^a dx^b = -e^{-2\delta} A dt^2 + A^{-1} dr^2 + r^2 d\phi^2, \quad (6)$$

where δ and A are functions of (t, r) . Next, we assume that the wave maps are corotational, that is

$$F = F(t, r), \quad \Phi = \phi. \quad (7)$$

Equation (3) reduces then to the single semilinear wave equation (hereafter, primes and dots denote derivatives with respect to r and t , respectively)

$$-e^\delta (e^\delta A^{-1} \dot{F})' + \frac{e^\delta}{r} (r e^{-\delta} A F')' = \frac{\sin(2F)}{2r^2}, \quad (8)$$

and the Einstein equations become

$$\dot{A} = -\alpha r A \dot{F} F', \quad (9)$$

$$\delta' = -2\Lambda r - \alpha r (F'^2 + A^{-2} e^{2\delta} \dot{F}^2), \quad (10)$$

$$A' = -\alpha r \left(A F'^2 + A^{-1} e^{2\delta} \dot{F}^2 + 2 \frac{\sin^2 F}{r^2} \right). \quad (11)$$

The studies of the initial value problem for this system in the case of zero cosmological constant, performed first in the flat spacetime ($\alpha = 0$) [2] and recently also for $\alpha > 0$ [5], showed that the scale-free static solution plays a crucial role in the process of singularity formation, namely singularities form via a static solution shrinking adiabatically to zero size. In fact, Struwe showed that for equivariant wave maps in the flat spacetime singularities must form in this way [6], in other words non-existence of a nontrivial static solution implies global regularity. Thus, it seems interesting to see how the inclusion of a negative cosmological constant affects the structure of static solutions.

2 Static solutions for $\Lambda = 0$

Before looking at the static solutions of equations (8)-(11) with $\Lambda < 0$, in this section we review some well known facts about static solutions for $\Lambda = 0$. We first consider the case $\alpha = 0$ which corresponds to the flat spacetime $A = 1, \delta = 0$ so equation (8) reduces to

$$\frac{1}{r} (r F')' = \frac{\sin(2F)}{2r^2}. \quad (12)$$

The trivial constant solutions of (12) are $F = 0$ and $F = \pi$; geometrically these are maps into the north and the south pole of S^2 , respectively. The energy of these maps

$$E(F) = \pi \int_0^\infty \left(F'^2 + \frac{\sin^2 F}{r^2} \right) r dr \quad (13)$$

attains the global minimum $E = 0$. Note that the requirement that energy be finite imposes a boundary condition at spatial infinity $F(\infty) = k\pi$ ($k = 0, 1, \dots$) which compactifies $\mathbb{R}^2$ into S^2 and thereby breaks the phase space into infinitely many disconnected topological sectors labelled by the degree k of the map $S^2 \rightarrow S^2$.

The fact that equation (12) is scale invariant does not exclude nontrivial regular solutions with finite energy (Derrick's argument is not applicable) and, in fact, such solutions are well-known both in the mathematical literature as harmonic maps from $\mathbb{R}^2$ into S^2 and in the physics literature as instantons in the two-dimensional euclidean sigma model. One way to derive them is to use the Bogomol'nyi identity

$$E(F) = \pi \int_0^\infty \left(F'^2 + \frac{\sin^2 F}{r^2} \right) r dr = \pi \int_0^\infty \left(\sqrt{r} F' - \frac{\sin F}{\sqrt{r}} \right)^2 dr - 2\pi \cos F \Big|_0^\infty. \quad (14)$$

It follows from (14) that in the topological sector $k = 1$ the energy attains the minimum, $E = 4\pi$, on the solution of the first order equation $rF' = \sin F$, which is

$$F_S(r) = 2 \arctan(r/\lambda), \quad (15)$$

where λ is a nonzero constant. This solution is a well-known harmonic map from R^2 to S^2 . We remark in passing that this solution can be alternatively obtained in an elegant geometric way by composing the identity map between two-spheres with the inverse of stereographic projection

It has been known for long that the solution (15) persists if one couples gravity with zero cosmological constant [1]. To see this let us consider equations (8)-(11) and assume that the fields are time independent. We get

$$\frac{1}{r} e^\delta (A e^{-\delta} r F')' = \frac{\sin(2F)}{2r^2}, \quad (16)$$

and (assuming that $\Lambda = 0$)

$$\delta' = -\alpha r F'^2, \quad (17)$$

$$A' = -\alpha r \left(A F'^2 + \frac{\sin^2 F}{r^2} \right). \quad (18)$$

For regular solutions the boundary conditions at $r = 0$ are

$$A(0) = 1, \quad \delta(0) = 0, \quad F(0) = 0, \quad F'(0) = b, \quad (19)$$

where b is a free parameter. We want a finite energy degree-one solution so we require that $A(r)$ and $\delta(r)$ tend to constants at infinity and $F(\infty) = \pi$. Such a solution can be found explicitly as follows. Let $B = \exp(-2\delta)A$. Then, from (17) and (18) we obtain

$$B' = \alpha r \left(AF'^2 - \frac{\sin^2 F}{r^2} \right) e^{-2\delta} \quad (20)$$

Using the boundary conditions (19), this implies that $B(r) \equiv 1$, hence $A = \exp(2\delta)$. Substituting this into (16) we get

$$\frac{1}{r} e^\delta (r e^\delta F')' = \frac{\sin(2F)}{2r^2}. \quad (21)$$

Using the new coordinate ρ defined by

$$r e^\delta \frac{d}{dr} = \rho \frac{d}{d\rho}, \quad (22)$$

one can rewrite equation (21) in the form of the flat space equation (12)

$$\frac{1}{\rho} \frac{d}{d\rho} \left(\rho \frac{dF}{d\rho} \right) = \frac{\sin(2F)}{2\rho^2}, \quad (23)$$

which, as we showed above, is solved by $F_S(\rho) = 2 \arctan(\rho/\lambda)$. Inserting this solution into equation (17) and integrating we get

$$e^\delta = 1 - \frac{2\alpha\rho^2}{\lambda^2 + \rho^2}. \quad (24)$$

This yields the metric

$$ds^2 = -dt^2 + (\lambda^2 + \rho^2)^{-2\alpha} (d\rho^2 + \rho^2 d\phi^2) \quad (25)$$

which has the deficit angle equal to $4\alpha\pi$, hence this solution exists only for $\alpha < 1/2$.

3 Non-existence of static solutions for $\Lambda < 0$

For a nonvanishing cosmological constant, the static equations (16) and (17) do not change while equation (18) picks up an additional term

$$A' = -2\Lambda r - \alpha r \left(AF'^2 + \frac{\sin^2 F}{r^2} \right). \quad (26)$$

Assuming that $\Lambda < 0$, by rescaling, without loss of generality, we set hereafter $\Lambda = -1$. Using (17) we eliminate δ from (16) and get the following system

$$A' = 2r - \alpha r \left(AF'^2 + \frac{\sin^2 F}{r^2} \right), \quad (27)$$

$$F'' + \frac{1}{r}F' + \frac{2r^2 - \alpha \sin^2 F}{Ar}F' = \frac{\sin(2F)}{2Ar^2}. \quad (28)$$

The boundary conditions at $r = 0$ are

$$F(r) \sim br, \quad A(r) \sim 1 - (\alpha b^2 - 1)r^2. \quad (29)$$

We want a solution for which $A \sim r^2$ at infinity and $F(\infty) = \pi$. Such a solution was claimed to exist and constructed numerically in [3]. We shall show now that this claim was erroneous.

Let us define a function

$$H = \cos^2 F + r^2 A F'^2. \quad (30)$$

We claim that $H(r)$ is monotone decreasing. To prove this, using the field equations, we compute

$$H' = -2r^3 F'^2 - \alpha r^3 A F'^4 + \alpha r \sin^2 F F'^2. \quad (31)$$

It is convenient to rewrite equation (31) in the form

$$H' = -r F'^2 G(r), \quad G(r) = 2r^2 - \alpha r^2 A F'^2 + \alpha \sin^2 F. \quad (32)$$

From the boundary conditions (29) $G(r) \sim 2r^2 > 0$ near $r = 0$, independent of b and α . Now we shall show that $G(r) = 0$ implies $G'(r) \geq 0$. To this end we compute

$$G'(r) = -2\alpha r^3 F'^2 - \alpha^2 r^3 A F'^4 + 4r + \alpha^2 r \sin^2 F F'^2. \quad (33)$$

To evaluate G' when $G = 0$ we solve $G = 0$ for $\sin^2 F$ and substitute that value into equation (33). We get

$$G' \Big|_{G=0} = 4r. \quad (34)$$

Thus $G'(r) > 0$ for $r > 0$, and therefore $H'(r) \leq 0$ for all r . Since $H(0) = 1$, this implies that $H(r) < 1$ for all $r > 0$. This excludes existence of a solution having $\lim_{r \rightarrow \infty} F(r) = \pi$ because that would mean $\lim_{r \rightarrow \infty} H(r) \geq 1$.

In view of Struwe's result mentioned above, the nonexistence of a static nonconstant soliton suggests (but by no means proves) that the negative cosmological constant might act as a cosmic censor in this model.

Acknowledgement

PB thanks Lars Andersson for discussions. The research of PB was supported in part by the KBN grant 2 P03B 006 23 and the FWF grant P15738.

References

- [1] A. Comtet and G. W. Gibbons, Nucl. Phys. **B299**, 719 (1988).
- [2] P. Bizoń, T. Chmaj, and Z. Tabor, Nonlinearity **14**, 1041 (2001).
- [3] Y. Kim and S.-H. Moon, Phys. Rev. D **58**, 105013 (1998).
- [4] G. Gibbons, M. E. Ortiz, and F. Ruiz Ruiz, Phys. Rev. D **39**, 1546 (1989).
- [5] T. Chmaj and S. J. Szybka, in preparation.
- [6] M. Struwe, Comm. Pure Appl. Math. **56**, 815 (2003).